

This article was downloaded by: [195.39.66.118]

On: 22 August 2012, At: 08:02

Publisher: Routledge

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Psychology & Health

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/gpsh20>

EHPS 2012 abstracts

Version of record first published: 01 Aug 2012

To cite this article: (2012): EHPS 2012 abstracts, Psychology & Health, 27:sup1, 1-357

To link to this article: <http://dx.doi.org/10.1080/08870446.2012.707817>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.tandfonline.com/page/terms-and-conditions>

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae, and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand, or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

Motivations for Aesthetic Plastic Surgery

P. Pires¹

¹Universidade Autonoma de Lisboa, Lisboa, Portugal

After bibliographic study we applied 17 semi structured interviews from which two sub categories arise: a) Physical motivations and b) Psychological motivations. Starting in the sense units we formulated a group of affirmations that referred to each one of the summoned categories and sub categories that in their group composed the first version of the questionnaire. This first version was presented to a group of 7 judges. The scale was then applied to a sample of 30 patients in two separate moments separated by three weeks (test-retest) in order to perceive the constancy of the results and the stability of the construct which was meant to measure. The scale was then applied to a sample of 122 patients of the aesthetical surgery appointments with the goal of scale validation which results (four factors that explain 63.3% of total variance) and internal consistence ($\alpha = 0.938$) guarantee its adequacy.